

# Activity concentration of $^{222}\text{Rn}$ gas, $^{226}\text{Ra}$ , $^{232}\text{Th}$ and $^{40}\text{K}$ in crops and soil taken from Safwan Granges using active, passive and gamma spectroscopy techniques

JABBAR H. JEBUR AND ABDUL RIDHA H. SUBBER\*

*Physics department, college of education of pure sciences, University of Basrah, Basrah, Iraq*

\*Email; [dabrhs@hotmail.com](mailto:dabrhs@hotmail.com)

## ABSTRACT

It is widely expected that fertilizer has great effect on the soil crops, especially when it contains a radioactive elements. The radioactive content present in crops will increase the number of deferent type of cancer cases in consumers and will explain the importance of this study. In this work we used the passive method (can technique) to measure radon concentration and gamma ray spectrometer for gamma ray concentrations measurement type NaI(Tl) 3"x3". The samples show an average radon concentration of  $48.58 \text{ Bq/m}^3$ , area radon exhalation rate of  $0.091 \text{ Bq.m}^{-2}.\text{h}^{-1}$  and mass exhalation rate of  $0.007 \text{ Bq.kg}^{-1}.\text{h}^{-1}$ . The average specific gamma activity concentration for  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$ ,  $^{238}\text{U}$  and  $^{40}\text{K}$  were found to be 37.94, 27.88, 13.2 and 388.0 Bq/kg respectively. The gamma hazard indices  $R_{\text{eq}}$ ,  $H_{\text{ex}}$ ,  $H_{\text{in}}$ ,  $D$ ,  $E_{\text{out}}$  were found to be 105.58 Bq/kg, 0.285, 0.383, 49.35 nGy/h, 0.06 mSv/y respectively. The concentrations of radionuclides found in this study are nominal and do not pose any health hazard.

**Keywords:** *radon, passive, active, radium, gamma spectroscopy.*

## 1. INTRODUCTION


The transfer of metals and radioactive element through the planets to humans is a serious health hazard. It mostly causes lung, stomach, kidney, bladder cancers as well as leukemia [1]. To increase the crop production and to improve soil nutrient properties, farmers used different categories of fertilizers which are essential in this time. Most of the materials contain  $^{238}\text{U}$  and  $^{232}\text{Th}$  are radon gas emitters, which is a progeny of Radium, which in turn is derived from Uranium decay. The natural radioactivity arises mainly from  $^{40}\text{K}$  and nuclide from the  $^{238}\text{U}$  and  $^{232}\text{Th}$  series and their decay product [2]. A combination of soil and fertilizer is essential for understanding changes in the background radiation [3]. The radionuclide (Uranium, Thorium, Radium and Potassium) can be transferred from soil mixed with fertilizer to roots and accumulate in various part of plant like leaves grain and stem [4, 5]. These nuclei

have a half- life comparable to the edge of the earth and enhanced 87% of the total radiation dose received, the rest are manmade [6,7]. The fertilized soil is considered as the main source of continuous radiation exposure to human and acts as a medium of migration of radionuclide to the biological system.

The aim of the present work is to investigate the activity concentration of NORM in fertilized soil and crops in Safwan Granges, using radon gas detection and gamma ray spectroscopy.

## 2. STUDY AREA

The study area falls in Safwan region near Kuwait border as shown in Figure 1. Safwan is lying just south-western part of Basrah Governorate, southern Al-Zubiar district, is enclosed by salt water on eastern side by Khor Al-Zubiar channel between longitudes  $47^{\circ} 52'$  and  $48^{\circ} 30'$  E, and latitude circles  $30^{\circ} 10'$  and  $29^{\circ} 50'$  N, with an area of 24 km<sup>2</sup> south-west, near border of Kuwait, in which sediments are made up alluvial fan and sandy Dibddibba deposits. Dibddibba aquifer has acquired its name from Dibddibba Formation which is composed of unconsolidated deposits, i.e. alluvial, often represented by the most common exploited aquifer. Soil of the area comprises essentially of ill-sorted gravely sand, with gypcrete top cover. The size of the gravels varies from coarse gravels (5 - 20 cm), around the apex of the fan to fine gravels and pebbles (2 - 5 cm), in the peripheral parts. The thickness of the gypcrete varies from (0.5 - 1.5)m. Geographically; the surficial fine-grained soil occupies about (50%) of Basrah area. Most cities of Basrah Governorate are located on this type of soil. These soils have a range of depth (7-15) m of depth in the area. The area contains many granges used fertilizers for crops growth [8-10].



**Fig.1. Area of study with samples marking (from Google earth).**

### **3. MATERIALS AND METHODS**

#### **3.1. Radon measurements**

##### **3.1.1 Passive technique**

Twenty soil and crop samples were collected from different locations in granges of tomatoes in Safwan region near the border of Kuwait. The samples name and number are listed in Table 1.

**Table 1 The samples name and number of crops and soil**

<b>Samplenumber</b>	<b>Sample ID</b>
1	SOIL-1
2	TOMATO-1
3	SOIL-2
4	OPTION ATROSY
5	SOIL-3
6	EGGPLANT
7	SOIL-4
8	TOMATO-2
9	SOIL-5
10	TOMATO-3
11	SOIL-6
12	TOMATO-4
13	SOIL-7
14	OKRA
15	SOIL-8
16	QRNABIT BROCCOIL
17	SOIL-9
18	TOMATO-5
19	SOIL-10
20	TOMATO-6

The can technique was adopted for estimation the radon concentration and exhalation rates for crops and soil samples [11]. The samples were dried in a hot air oven at a temperature 110°C for 24 h and then crushed and sieved for uniform emanation of radon. Small masses (0.22 kg) were put inside hard plastic containers of 7.5 cm diameter and 30 cm height. In this study we used CR-39 track detectors to measure the track density. The detector 1.5 cm x 1.5 cm areas was fixed on the top inside of the containers using double side adhesive tape. A semi-permeable membrane was fitted in between the detector and sample, which allow only radon to diffuse through. The exposing time was 90 days, and then the detectors were removed and etched with 6.25N NaOH at 70°C during 7h. After etching, the detectors were

washed in distilled water and then dried in air. The track density, due to alpha particles emitted by samples, was determined by using optical microscope (400X). The radon concentration was calculated using the following expression [12,13];

$$A_{Rn} \left( \frac{Bq}{m^3} \right) = \frac{\rho}{TK} \quad (1)$$

Where T is the exposure time in day,  $\rho$  is track density in T/cm<sup>2</sup> and K is the calibration factor, which used as K= 0.2857±0.0143 Tr/cm<sup>2</sup>.d per Bq/m<sup>3</sup> measured from previous calibration experiment [11].

At the equilibrium state, the radon flux (exhalation) from each sample inside the can is written as [14, 15];

$$E_A = \frac{A_{Rn}VT\lambda/S}{T+(e^{-\lambda T}-1)/\lambda} \quad (2)$$

where E<sub>x</sub> is exhalation rate in unit Bq.m<sup>-2</sup>.d<sup>-1</sup>, A<sub>Rn</sub> is radon concentration measured by the detector in unit Bq.m<sup>-3</sup>,  $\lambda$  is radon decay constant, T is the exposure time, V is the volume of the can and S is the surface area of sample in the can.

The radon exhalation rate in terms of mass is calculated from the relation;

$$E_M = \frac{A_{Rn}VT\lambda/M}{T+(e^{-\lambda T}-1)/\lambda} \quad (3)$$

where E<sub>M</sub> expressed in Bq.kg<sup>-1</sup>.h<sup>-1</sup> and M in kg is the mass of the sample.


The effective radium content calculated from the relation [16];

$$A_{Ra} = \frac{\rho hS}{KMT} \quad (4)$$

where  $\rho$  is track density recorder, h distance between the detector and sample, S surface area of sample, T is exposure time and K is the calibration factor.

### 3.1.2. Active technique

4. RAD7 (DurrIDGE Company) is a highly versatile electronic instrument used for radon detection that can measure radon concentrations in real time. For the present purposes the instrument was used to measure radon concentration emanated from soil. The soil sample was loaded into 30 cm x 7.5 cm can used as an emanation cylinder. The cylinder connected online with RAD7 system as shown in Figure 2. The height of the active volume of the container is 25 cm, to insure <sup>222</sup>Rn detection only. The alpha RAD7 detector was operated in grab mode for one day protocol, with cycle 0.5 h and recycle 48. The removable lid was equipped with two gas-tight tubes, one used to pump air contains radon gas in RAD7 chamber (ZnS) and the other used to pump fresh air to the container. The system is a closed loop in which the air circulates continuously. The concentration of radon emanated from each sample inside the emanation chamber was allowed to build up with time and it was measured in 0.5h cycle for an average time of 24h. The output data are produced by computer software (Capture 1.2.0) was supplied by DURRIDGE Company.



**Fig. 2. Block diagram of RAD7 instrument online with sample dosimeter for active radon measurement.**

#### **4.1. Gamma ray measurements**

Samples of crops and soil were heated in the oven at 110°C for 24h to remove moisture, grinded to fine powder, put inside Marinalli beakers and then stored for 30days to allow the equilibrium between  $^{226}\text{Ra}$  and  $^{222}\text{Rn}$ . The activity concentration of  $^{226}\text{Ra}$ ,  $^{228}\text{Ra}$ ,  $^{238}\text{U}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  was estimated from the gamma spectrum using Na(Tl) detector 3x3 inch with a 1024 channel computer analyzer USX supplied by Spectrum Technique Company. The detector was employed with lead shielding, 4 cm thickness, which reduced the background. The detector was calibrated using standard sources of  $^{57}\text{Co}$  (peak 122 keV),  $^{137}\text{Cs}$  (peak 662 keV) and  $^{60}\text{Co}$  (peaks 1173, 1333 keV). The detector resolution is about 8% at 662 keV of  $^{137}\text{Cs}$ . The efficiency calibration was achieved using eight standard sources include the calibration sources. The system was running freely, for 12 h live time, to evaluate the background spectrum. The Marinalli beaker contains sample was placed over the detector for counting.


After measuring the net count (area under the peak) for each peak, the activity concentration for each environmental isotope calculated from [17];

$$A = \frac{\text{Net count}}{\varepsilon \times I_{\gamma} \times M \times t} \quad (5)$$

where  $\varepsilon$  is absolute gamma peak efficiency of the detector at this particular gamma-ray energy,  $I_{\gamma}$  decay intensity for the specific energy peak (including the decay branching ratio

information), M the mass of the sample in kg and t is the counting time of the measurement in second.

For calculation of the specific activity concentration for each Normal Origin Radioactive Material (NORM), or Technical Enhanced Normal Origin Radioactive Materials (TENORM), one has to recognize the belongcity of each peak according to gamma decay of each isotope. For calculation of the specific activity concentration for each environmental (NORM), or (TENORM), one has to recognized the belong city of each peak according to gamma decay of each isotope. The activity concentration of  $^{226}\text{Ra}$ , is calculated from the weighted average concentration of gamma ray lines; 295 keV(19.2%), 352 keV (37.1%), 609 keV ( 46.1%), 1120 keV (15%) and 1760 keV (15.4%). The peak of 186 keV assumed to be from  $^{235}\text{U}$  since it has slight effect on the total concentration after subtracting the background, 42.8% for Ra and the rest for  $^{235}\text{U}$ . The determination of existence of  $^{232}\text{Th}$  is achieved by the average of lines; 338 keV (12%), 969 keV (17%). The case of  $^{238}\text{U}$  is recognized by 1001 keV (83%), 766 keV (29%) and 2204 keV (5%). For  $^{40}\text{K}$ , this directly determined using 1460 keV (10%) peak [18]. Sample of environmental gamma ray spectrum in 512 is shown in figure 3.



**Fig. 3. Gamma ray spectrum of soil sample measured by NaI.**

Radium equivalent activity ( $Ra_{eq}$ ) is used to assess the hazards associated with materials that contain  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  in  $\text{Bq kg}^{-1}$ , which is, determined by assuming that  $370 \text{ Bq kg}^{-1}$  of  $^{226}\text{Ra}$  or  $260 \text{ Bq kg}^{-1}$  of  $^{232}\text{Th}$  or  $4810 \text{ Bq kg}^{-1}$  of  $^{40}\text{K}$  produce the same  $\gamma$  dose rate. The  $Ra_{eq}$  of a sample in ( $\text{Bq kg}^{-1}$ ) can be achieved using the following relation [19];

$$Ra_{eq} = (A_{Ra}) + (A_{Th} \times 1.43) + (A_K \times 0.077) \quad (6)$$

The published maximal permissible  $Ra_{eq}$  is 370 Bq kg<sup>-1</sup>.

The external and internal hazard indices are an evaluation of the hazard of the natural gamma radiation. The prime objective of this index is to limit the radiation dose to the admissible permissible dose equivalent limit around 1mSv<sup>-1</sup>. In order to evaluate this index, one can use the following relations[20]

$$H_{ex} = (A_{Ra}/370) + (A_{Th}/259) + (A_K/4810) \quad (7)$$

$$H_{in} = (A_{Ra}/185) + (A_{Th}/259) + (A_K/4810) \quad (8)$$

In order to estimate the annual effective dose rate in air, the conversion coefficient from absorbed dose in air to effective dose received by an adult must be considered. This value is published in UNSCEAR 2000 and UNSCEAR 1993, to be 0.7 SvGy<sup>-1</sup> for environmental exposure to gamma rays of moderate energy. The outdoor occupancy factor is about 0.2 [18]. The annual effective dose equivalent is given by the following equation [19];

$$AEDE_{out}(mSv/y) = D(nGy/h) \times 8760(h/y) \times 0.2 \times 0.7(Sv/Gy) \times 10^{-6}(9)$$

$$\text{where } D\left(\frac{nGy}{h}\right) = 0.0417A_K + 0.462A_{Ra} + 0.606A_{Th}$$

The world average annual effective dose equivalent (AEDE) from outdoor or indoor terrestrial gamma radiation only is 0.560 mSv/year [UNSCEAR].

## 4. RESULTS AND DISCUSSION

### 4.1. Radon results

Table 2 shows the results of radon concentrations measured by passive (using equation 1) and active technique by RAD7.

**Table 2 Radon concentrations in samples collected from Safwan granges measured by passive and active methods.**

Sample No	Radon conc. in Bq/m <sup>3</sup> by RAD7	Radon conc. in Bq/m <sup>3</sup> by CR-39
1	94.4±7.0	106.4±35.3
2	28.5±4.2	22.0±8.6
3	60.8±5.9	66.0±22.6
4	29.9±3.6	20.8±8.3
5	45.6±4.7	51.4±17.9
6	47.5±5.2	39.1±14.1
7	46.0±6.2	57.5±19.9
8	21.8±3.6	24.5±9.4
9	37.2±5.2	25.7±9.8
10	36.0±4.8	18.3±7.4
11	137.0±25.0	97.8±32.6
12	38.5±7.2	31.8±11.7
13	38.8±5.4	36.7±13.3

14	22.4±4.2	20.8±8.3
15	52.0±6.2	50.1±17.6
16	22.0±3.8	20.8±8.3
17	54.4±5.4	51.4±17.9
18	34.3±3.7	31.8±11.7
19	54.6±5.9	80.7±27.3
20	25.8±3.4	23.2±9.1

Figure 4 shows the correlation coefficient between passive and active techniques this correlation equal ( $R^2=0.95$ ), which means that, the correlation is excellent.




Fig.4. Correlation between active (using RAD7) and passive (using CR-39) techniques.

The maximum value of radon concentration in table 2; is  $137.0 \pm 25.0 \text{ Bq/m}^3$  in sample number 11 (Soil-6), measured by RAD7 and minimum value is  $18.3 \pm 7.4 \text{ Bq/m}^3$  in sample number 10 (Tomatoes-3), measured by CR39. The arithmetic mean value is  $48.58 \text{ Bq/m}^3$ , which is very low in comparison with the international recommended values. Table 3, presented the radon area exhalation rate, mass exhalation rate and effective mass concentration of radium which decays to produce such parameters by using equations 2, 3 and 4. The results show the range of  $E_A=0.037$  to  $0.214 \text{ Bq/m}^2 \cdot \text{h}$  and the average value is  $0.091 \text{ Bq/m}^2 \cdot \text{h}$ . These results are consistent with many researches works, for example Chauhan et al. [1] worked on tobacco leaves and found that the average exhalation rate  $0.115 \text{ Bq/m}^2 \cdot \text{h}$  and the work of Mahur et al. [21] measured the average exhalation rate from soil as  $0.119 \text{ Bq/m}^2 \cdot \text{h}$ . The effective radium in the samples is in the range from  $0.098$  to  $0.57 \text{ Bq/kg}$ , with arithmetic average  $0.244 \text{ Bq/kg}$ .

**Table 3 Area exhalation rate, mass exhalation rate and the effective radium mass.**



Sample No	$E_A$ in $Bq.m^{-2}.h^{-1}$	$E_m$ in $Bq.m^{-2}.h^{-1}$	$A_{Ra}$ in $Bq/kg$
1	0.214±0.073	0.017±0.006	0.570±0.036
2	0.044±0.018	0.004±0.001	0.118±0.010
3	0.133±0.046	0.011±0.004	0.354±0.024
4	0.042±0.017	0.003±0.001	0.111±0.009
5	0.103±0.037	0.008±0.003	0.275±0.019
6	0.079±0.029	0.006±0.002	0.210±0.015
7	0.115±0.041	0.009±0.003	0.308±0.021
8	0.049±0.019	0.004±0.002	0.131±0.010
9	0.052±0.020	0.004±0.002	0.138±0.011
10	0.037±0.015	0.003±0.001	0.098±0.008
11	0.197±0.067	0.016±0.005	0.524±0.034
12	0.064±0.024	0.005±0.002	0.170±0.013
13	0.074±0.027	0.006±0.002	0.197±0.014
14	0.042±0.017	0.003±0.001	0.111±0.009
15	0.101±0.036	0.008±0.003	0.269±0.019
16	0.042±0.017	0.003±0.001	0.111±0.009
17	0.103±0.037	0.008±0.003	0.275±0.019
18	0.064±0.024	0.005±0.002	0.170±0.013
19	0.162±0.056	0.013±0.004	0.432±0.028
20	0.047±0.019	0.004±0.001	0.124±0.010
<b>Max</b>	0.214±0.073	0.017±0.006	0.570±0.036
<b>Min</b>	0.037±0.015	0.003±0.001	0.098±0.008
<b>Aver.</b>	<b>0.091</b>	<b>0.007</b>	0.244

#### 4.1. Gamma ray results


Table 4 shows the specific natural activity concentrations of  $^{226}Ra$ ,  $^{228}Ra$ ,  $^{232}Th$ ,  $^{238}U$  and  $^{40}K$  existing in samples collected from selected areas in Safwan granges. For  $^{226}Ra$ , the specific activity ranged from 21.39±6.31 to 68.78±6.31 Bq/kg with an average 37.94±4.92 Bq/kg. For  $^{232}Th$ , the specific activity ranged from 0.07±0.06 to 76.53±5.33 Bq/kg. For  $^{238}U$ , specific activity ranged from 0.04±0.01 to 64.81±17.1 Bq/kg with an average 13.28±3.17 Bq/kg and for  $^{40}K$  it ranged from 43.8±18.0 to 1000±19 Bq/kg with an average 388.3±5.7 Bq/kg. Therefore, the results clearly indicate that all the radionuclides detected were in nominal concentration.

**Table 4 The specific activity concentration for natural radionuclides**

S. N	Ra-226 Bq/kg	Th-232 Bq/kg	Ra-228 Bq/kg	K-40 Bq/kg	U-238 Bq/kg
1	68.78±6.31	40.6±2.94	25.58±1.49	261.2±1.1	0.51±0.01
2	28.36±2.86	11.16±0.89	35.68±2.35	699.1±8.2	2.19±0.09
3	50.49±3.49	45.85±3.29	66.82±3.90	285.0±1.3	0.45±0.02
4	28.5±3.31	14.96±1.30	33.69±2.47	937.0±17.7	2.93±0.14
5	39.54±2.71	24.86±2.36	26.10±1.53	272.25±1.2	0.04±0.02
6	31.72±5.59	61.57±5.22	58.3±28.54	1000 ±19	5.74±0.20


7	42.37±2.85	37.28±3.11	41.85±2.44	274.4±1.2	8.56±1.81
8	29.7±4.43	3.16±1.36	0.06±0.04	879.0±2.0	8.43±0.30
9	23.11±1.63	29.75±2.19	44.89±2.59	239.9±1.0	0.85±0.01
10	21.39±3.65	0.60±0.88	23.67±1.56	556.0±6.5	10.03±2.55
11	61.17±4.88	36.52±2.78	39.24±2.79	281.7±1.3	2.29±0.55
12	30.71±4.74	0.60±0.74	18.90±1.25	58.8±0.7	6.71±1.68
13	31.54±4.89	2.87±0.21	0.02±0.01	43.9±0.2	18.01±4.42
14	25.55±6.73	9.30±0.80	10.91±0.81	229.7±4.4	64.81±17.10
15	38.45±7.21	30.19±2.13	0.02±0.01	89.1±0.4	39.26±9.64
16	27.28±5.92	0.07±0.06	2.31±0.21	110.5±1.5	0.36±0.19
17	41.29±8.86	76.53±5.33	3.11±0.18	44.7±1.8	0.17±0.02
18	33.26±6.56	28.12±2.51	12.08±7.90	215.0±4.1	48.86±12.70
19	59.74±7.1	64.49±4.6	39.28±6.03	217.8±2.9	3.66±0.92
20	31.66±4.04	18.38±1.78	51.47±32.81	802.7±15.5	3.38±0.15
<b>Max.</b>	<b>68.78±6.31</b>	<b>76.53±5.33</b>	<b>66.82±3.28</b>	<b>1000±19</b>	<b>64.81±17.10</b>
<b>Min.</b>	<b>21.39±3.65</b>	<b>0.07±0.06</b>	<b>0.02±0.01</b>	<b>43.8±18.0</b>	<b>0.04±0.01</b>
<b>Ave.</b>	<b>37.94±4.92</b>	<b>27.88±2.62</b>	<b>27.31±5.96</b>	<b>388.3±5.7</b>	<b>13.28±3.1</b>

Figure 5 shows the correlation between total radium concentrations measured by gamma ray spectroscopy and radon measured by CR39 track detectors or exhalation rate. There looks to be strong and positive correlation between them (R=0.98).



**Fig.5. Correlation between Ra-226 measured by NaI and Rn-222 measured by CR-39**

Figure 6 shows the correlation between effective radium concentrations measured by CR-39 track detectors and total radium activity concentrations measured by gamma ray spectroscopy. The correlation looks positively strong, with correlation factor  $R=0.98$ .



**Fig.6. Correlation between total  $^{226}\text{Ra}$  activity concentration measured by NaI and effective  $^{226}\text{Ra}$  measured by CR-39 track detector.**

A detailed analysis of the results in table 4, indicates that there are no correlations between radioactive elements belong to different series. Figures 7, 8 and 9 were introduced to check this correlation.




Fig. 7. Correlation between  $^{232}\text{Th}$  and  $^{238}\text{U}$  activity concentrations in the samples





Fig. 8. Correlation between  $^{228}\text{Ra}$  and  $^{226}\text{Ra}$  activity concentrations in the samples



**Fig.9. Correlation between  $^{226}\text{Ra}$  and  $^{40}\text{K}$  activity concentrations in the samples.**

The radium equivalent activity, external and internal hazard index and annual effective dose calculated using equations (6) to (9) are presented in Table 5.

The results revealed that the measured activity levels from natural occurring radioactive in the investigated soil and crops samples are comparable to the corresponding worldwide level. The calculated result of radium equivalent activity ranged from  $35.85 \pm 6.11$  Bq/kg to  $196.77 \pm 14.50$  Bq/kg with a mean value of  $105.58 \pm 13.60$  Bq/kg, and is lower than 370 Bq/kg reported by UNCEAR2000. The average values of  $H_{\text{ex}}=0.282$  and  $H_{\text{in}}=0.383$  are less than 1 as recommended.

**Table 5 Values of radiation hazard parameters for samples.**

Sample No	$Ra_{\text{eq}}$ Bq/kg	$H_{\text{ex}}$	$H_{\text{in}}$	DnGy/h	$AEDE_{\text{out}}$ mSv/y
1	$146.94 \pm 10.61$	0.396	0.582	65.892	0.080
2	$98.22 \pm 4.76$	0.265	0.341	49.162	0.060
3	$137.99 \pm 8.29$	0.372	0.509	62.378	0.076
4	$122.11 \pm 6.52$	0.329	0.406	61.822	0.075
5	$96.05 \pm 6.17$	0.259	0.366	44.077	0.053
6	$196.77 \pm 14.50$	0.531	0.617	94.910	0.115
7	$116.82 \pm 7.39$	0.315	0.430	53.120	0.064

8	101.91±7.91	0.275	0.355	52.450	0.064
9	84.12±4.83	0.227	0.289	38.720	0.047
10	65.06±5.40	0.175	0.233	33.410	0.040
11	135.07±8.95	0.364	0.530	60.981	0.074
12	36.08±5.85	0.097	0.180	16.010	0.019
13	39.02±5.19	0.105	0.190	17.144	0.020
14	56.52±8.21	0.152	0.221	26.578	0.032
15	88.48±10.28	0.239	0.342	39.059	0.047
16	35.85±6.11	0.096	0.170	16.425	0.020
17	154.17±16.49	0.416	0.527	67.232	0.082
18	90.02±10.47	0.243	0.333	40.967	0.050
19	168.73±13.75	0.455	0.617	75.052	0.091
20	119.75±7.80	0.323	0.408	59.485	0.072
<b>Max.</b>	<b>196.77±14.50</b>	<b>0.531</b>	<b>0.617</b>	<b>94.91</b>	<b>0.115</b>
<b>Min.</b>	<b>35.85±6.11</b>	<b>0.096</b>	<b>0.17</b>	<b>16.01</b>	<b>0.019</b>
<b>Ave.</b>	<b>105.58±13.6</b>	<b>0.285</b>	<b>0.383</b>	<b>49.35</b>	<b>0.060</b>

The observed gamma dose rate varies from 16.01 to 94.91 nGy/h with an average value of 49.35 nGy/h. This result is less than the reported value of UNSCEAR2000 [22] which is approximately 57nGy/h. The annual effective dose varies from 0.019 mSv/y to 0.115 mSv/y with the mean value of 0.06 mSv/y. The worldwide average effective dose, according to UNSCEAR2000 is 0.07 mSv/y, this means that, our results were in agreement with the worldwide average value. In other word, the fertilized soil and the crops are safe to deal with.

## 5. CONCLUSIONS

The present measurements concentrate on data analysis for radon concentration and gamma ray emission from soil and crops samples collected from Safwan granges near the border of Kuwait. The samples show acceptable radon gas concentration in fertilized soil and crops. Area radon exhalation and mass exhalation have been calculated and show low values. The activity concentration of radium-226, thorium-232 and potassium-40 were measured also for the same samples. The values were below the recommended limit by the ICRP as the maximum annual dose to member of public. The radium equivalent activity and hazard indices are found to be lower than the safe limit. The values of the annual observed effective

show that the radioactivity of the natural radionuclides found in the surveyed area is nominal and does not pose any potential health hazard to the farmers and occupiers.

## REFERENCES

1. Chauhan P., Chauhan R. P. Measurement of fertilizers induced radioactivity in tobacco plants and elemental analysis using ICAP-AES. *Radiation Measurements*, 2014;63: 6-11.
2. Al- Jaralla M. I., Abu-Jarad F., Rehman, F. Determination of radon exhalation rates from tiles using active and passive technique. *RadiationMeasurements*,2001; 34: 491-495.
3. El-Farrash A. H., Yousef, H. A , Hafez, A. F. Activity concentration of  $^{238}\text{U}$  and  $^{232}\text{Th}$  in some soil and fertilizer samples using passive and active technique. *Radiation Measurements*, 2012; 47: 644-648.
4. Juber J. H.,Subber Abdul R. Natural radioactivity of some local and imported fertilizer in Basrah Governorate/Iraq., Scholar research library, Archive of Physics Research, 2014; 5(5): 18-22.
5. Kant K., Upadhyay S. B., Sonkawade R. G. ,Chakarvarti S. K. Radiological risk assessment of use of phosphate fertilizers in soil. *Iran j. Radiat. Res.* 2006; 4(2): 63-70.
6. Hamideen M. S.,Sharaf J. Natural radioactivity investigation in soil samples obtained from phosphate hills in the Russaifa region, Jordan. *Radiation Physics and Chemistry*, 2012; 81: 1559-1562.
7. Al-Trabusly H. A., Khater A.E.M,Habbani F. I. Radioactivity level and radiological hazard indices at the Saudi coastline of the Gulf of Aqaba. *Radiation Physics and Chemistry*, 2011;80: 343-348.
8. Shareef Najed F., Mahdi Maher M., Studying of recent environments in Faw, Khor Al-Zubair and Um-Qaser areas, South western Arabian Gulf, Basrah, Iraq. *Journal of Basrah Researches –Sciences A*, 2015: 41(2):1-14.
9. Sissakian Varoujan K., Shihab Ahmad T., Al-Ansari Nadhir , Knutsson Sven, Al-Batin Alluvial Fan, Southern Iraq. *Engineering*, 2014, 6: 699-711.
10. Khedher W.M. and Khorshid S.Z., Determination of the saline water intrusion zone and its contamination with ground water in the Dibddiba aquifer using vertical electrical sounding technique at Basrah Governorate, southern Iraq. *Mesopot. J. Mar. Sci.*, 2013, 28(2): 93 – 108.
11. Jebur J. H.,Subber R. H. Natural Radon exhalation rate from fertilizer used in Basra Governorate/ Iraq., *International Journal of Physics and Research (IJPR)*. 2014; 4(6): 1-10.
12. Elzain A.A., Measurement of Radon-222 concentration levels in water in Sudan. *Advances in applied Science Research*, 2014; 5(2): 229-234.

13. Alzoubi F. Y., Al-Azzam K.M, Alqadi M. K., Al-khateeb H.M., Ababneh Z.Q., Ababneh A.M. Radon concentration levels in the historical city Jeresh, Jordan, *Radiation Measurements*, 2013; 49: 35-38.
14. Al-Ghamdi S. S. Seasonal and location dependence of indoor and soil radon concentration in two villages, Nigran region, Saudi Arabia, *Radiation Measurements*, 2014; 69:12-17.
15. Imme G., Catalano R., Mangano G, Morelli D. Radon exhalation measurements for environmental and geophysics study. *Radiation Physics and Chemistry*, 2014; 95:349-351.
16. Somogyi G., Hafez A., Hunyadi I., Toth-Szilagyi, Measurement of exhalation and diffusion parameters of radon in solids by plastic track detectors. *Nucl. Tracks Radiat. Meas.*, 1986; 12(1-6):701-704.
17. Alharbi W. R. Natural Radioactivity and Dose Assessment for Brands of Chemical and Organic Fertilizers Used in Saudi Arabia. *Journal of Modern Physics*, 2013;4:344-348.
18. Papachristodoulou C. A., Assimakopoulos P.A, Paronis N E., Loannides K. G. Use of HPGe  $\gamma$ -ray spectroscopy to assess the isotopic composition of uranium in soil. *Journal of Environmental Radioactivity* 2003; 64:195-203.
19. Aziz Ahmed Qureshi, Ishtiaq Ahmed Khan Jadoon, Ali Abbas Wajid, Ahsan Attique, Adil Masood, Muhammad Anees, Shahid Manzoor, Abdul Waheed, Aneela Tubassam, *Radiation Protection Dosimetry*. 2014;158(4):466–478, doi:10.1093/rpd/nct271
20. UNSCEAR, 2008, United Nation scientific Committee on Effect of Atomic Radiation, VII
21. Mahur A. K., Gupta M., Varshney R., Sonkwade R. G. , Verma K. D. Radon exhalation and gamma radioactivity levels in soil and radiation hazard assessment in the surrounding area of National Thermal Power Corporation, Dadri, U.P, India. *Radiation Measurements*, 2013; 50, 130- 135.
22. UNSCEAR, 2000, United Nation scientific Committee on Effect of Ionized Radiation, Vol. II.