

1 Original Research Article  
2 **Comparative Investigation of the Antibacterial**  
3 **and Antifungal Potentials of the Extracts of**  
4 **Watermelon (*Citrullus lanatus*) Rind and Seed**

5  
6  
7  
8 **ABSTRACT**  
9

**Aims:** To investigate the antibacterial and antifungal potentials of the crude ethanol and aqueous extracts of watermelon (*Citrullus lanatus*) rind and seed.

**Study design:** The fruit was purchased from a local market and identified in the Department of Plant Science and Biotechnology, Michael Okpara University of Agriculture Umudike, Nigeria by a taxonomist. The rind and seed samples were extracted using ethanol (95%) and water and the crude extracts screened for antibacterial and antifungal potentials.

**Place and Duration of Study:** Department of Biochemistry, Michael Okpara University of Agriculture Umudike, Nigeria and National Root Crop Research Institute Umudike, Nigeria, between August 2014 and November 2014.

**Methodology:** The activities of the ethanol and aqueous extracts of watermelon (*Citrullus lanatus*) rind (EER and AER) and seed (EES and AES) at a concentration of 100 mg/ml were investigated by well-diffusion method against ten pathogens.

**Results:** The EER exhibited the highest activity against the bacterium, *Escherichia coli* (6.0) and the fungus, *Candida albican* (4.0). The least activity was in the EES against the fungus, *Trichosporo bePELLI* (0) followed by EER and AES against the bacterium, *Staphylococcus aureus* (1.0). For the bacterial strains, the highest mean activity (4.0) and mean susceptibility (4.50), respectively were in the EER and *Escherichia coli* whereas the least mean activity (3.0) and mean susceptibility (1.50) were in AES and *Staphylococcus aureus*. For the fungal strains, the highest mean activity (2.5) and mean susceptibility (3.0), respectively were in the EER, AER, AES and *Candida albican* whereas the least mean activity (1.25) and mean susceptibility (1.25) were in EES and *Trichosporo bePELLI*. The activities however were higher in the EE than in the AE of the rind and seed, suggesting the preference to the EE for higher pharmacologic activity.

**Conclusion:** The extracts (EER, AER, EES and AES) exhibited activities against the tested strains, especially *Escherichia coli* and *Candida albicans*, hence a scientific basis for their exploitation in nutraceutical formulations and in ethnomedication. The study underscores the overriding potency of EER against the bacterium, *Escherichia coli* and the fungus, *Candida albicans*, and the non-susceptibility of the fungus, *Trichosporo bePELLI* to EES. Further works, however are needed to validate reliability.

10  
11 **Keywords:** Nutraceutical, pharmacologic, pathogens, bacterial strains, fungal strains

12  
13 **1. INTRODUCTION**

14 The search for plants and plant parts with pharmacologic activity is warranted because of the increasing emergence of  
15 hitherto unknown and drug resistance diseases [1,2,3] and the need for a scientific basis for ethnomedicinal use of plants  
16 and plant parts extracts [4,5]. The use of plant based drugs in traditional medicine within African and the world over has  
17 been reported [6]. In particular, the medicinal value of fruits associated with their natural bioactive phytochemical contents  
18 [7] tend to increase fruit consumption the world over. In Nigeria, the pulp of various fruits, but not the rind and seed, is  
19 usually consumed. This contributes to food wastes in the environment as is the case with watermelon (family

cucurbitaceae and specie *Citrullus lanatus*), a major fruit widely distributed in the tropics that serves as a thirst-quencher owing to its high (92 %) water content [8,9]. To prevent solid waste related hazards to the environment, effort should be made to increase the utilization of food wastes which requires studies on their properties for scientific basis.

Water extract of watermelon seed is used in traditional herbal medicine to cure catarrhal infection, fever, disorders of the bowel and urinary passage [10]. Studies on watermelon fruits were reported, but mainly on the juice/pulp [11,12] and a little on the peel/rind [13,14]. These warranted this study aimed at comparing the antibacterial and antifungal potentials of the crude ethanol and aqueous extracts of watermelon (*Citrullus lanatus*) rind and seed against these bacterial strains viz: *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Bacillus subtilis*, *Bacillus pumilus*, *Staphylococcus aureus* and fungal strains viz: *Candida albicans*, *Aspergillus niger*, *Penicillium chrysogenum* and *Trichosporo bePELLI*.

## 2. MATERIAL AND METHODS

### 2.1. Chemicals and reagents

All solvent (ethanol) and other chemicals used, including those used in the preparation of reagents, were of analytical grade and products of reputable companies. This study was carried out between August and November, 2014.

### 2.2. Test organisms

The test organisms (pure culture of bacteria and fungi strains) used were bought from HOSLAB company Umuahia, Abia State, Nigeria. These include six bacterial strains (*Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Bacillus subtilis* and *Bacillus pumilus*) and four fungal strains (*Aspergillus niger*, *Candida albicans*, *Trichosporo bePELLI* and *Penicillium Chrysogenum*).

### 2.3. Collection and preparation of samples

Watermelon fruits were bought from Onuimo market, in Imo State border/boundary with Abia State, Nigeria. It was identified as Charleston gray variety in the Department of Plant Science and Biotechnology, Michael Okpara University of Agriculture Umudike, Nigeria. The watermelon was thoroughly washed to remove sand particles after which it was sliced using a home choice European knife. The seeds were handpicked and washed off the pulp particles using clean water. The pulp was carefully scraped off to obtain the rind which was chopped into pieces with a chipping machine.

The rind chips and seeds were respectively weighed, using Satorious Digital Weighing Balance, Model BP210S, Germany. The rind (wet weight = 1900.7 g) and seed (wet weight = 1016.9 g) were separately spread on a foil and sundried for 3 days to obtain the corresponding dry weight for the rind (82.6 g) and seed (468.5 g). The respective dry weight samples were milled into powder using Arthur Thomas Laboratory Mill, Crypto model, USA, covered separately in a labeled white nylon and kept in the desiccators until used.

### 2.4. Preparation of ethanol extract

To 10 g each of ground *Citrullus lanatus* seed and rind in a 250 ml conical flasks were added 200 ml of ethanol and the content allowed to settle for 24 hrs. The filtrate of the extracts was obtained by separation of the suspension with a filter paper (Whatman filter paper No 1). The ethanol extracts were allowed to evaporate and then stored in an airtight conical flask until used.

### 2.5. Preparation of aqueous extract

The aqueous extracts of *Citrullus lanatus* samples were prepared by squeezing the sand-free specimen in triple distilled water. The resultant solution was filtered and dialyzed by using sigma dialysis membrane-500 (average flat width, 24.26 mm; average diameter, 14.3 mm, and approximate capacity, 1.61 mlcm<sup>-1</sup>) against D-glucose to remove the excess water. The supernatant so obtained was lyophilized (Labcono Freeze Dry System) and stored at 4 °C in a refrigerator until used.

### 2.6. Preparation of media

7 gram each of nutrient agar and potato dextrose agar was weighed out in a beaker and dissolved in 250 ml of distilled water. The respective solutions were separately stirred, covered with a foil, autoclaved for 3 hours at 212 °C, allowed to cool in the autoclave (after reducing the temperature to 0 °C) and separately poured into the Petri dishes with the different bacteria and fungi strains.

### 2.7. Antibacterial activity

Petri dishes with nutrient agar were inoculated with six different species of bacteria. *Citrullus* species extraction was sterilized by passing each through a 0.22: m Millipore GV filter paper (Millipore USA). Round paper dishes with a radius of 0.8 cm were dipped into each extract (at a concentration of 100 mg/ml) and placed in the center of the inoculated petri-dishes. Bacterial colonies were allowed to grow overnight (24 hours) at 37 °C, and the inhibition zone (activity) around the disc measured.

### 2.8. Antifungal activity

Petri dishes with potato dextrose agar were inoculated with three different species of fungus. *Citrullus* species extracts were sterilized by passing each through a 0.22: m Millipore GV filter paper (Millipore USA). Round paper dishes with a radius of 0.8cm were dipped into each extract (at a concentration of 100 mg/ml) and placed in the center on inoculated petri-dishes. Fungal colonies were allowed to grow 48 hours at 28 °C and the inhibition zone (activity) around the disc measured.

### 2.9. Data

Data were presented simple mean obtained from duplicate measurement of each sample.

80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106

**3. RESULTS AND DISCUSSION**

The pharmacologic activities of the crude ethanol extracts (EE) and aqueous extracts (AE) of watermelon (*Citrullus lanatus*) rind (EER and AER) and seed (EES and AES) at a concentration of 100 mg/ml were investigated by well-diffusion method against ten pathogens (six bacterial and four fungal strains). The EER exhibited the highest activity/inhibition zone diameter (mm) against the bacterium, *Esherichia coli* (6.0) and the fungus, *Candida albican* (4.0). The least activity was in the EES against the fungus, *Trichosporo bePELLi* (0) followed by EER and AES against the bacterium, *Staphylococcus aureus* (1.0) (Tables 1 and 2).

**Table 1: Activity/inhibition zone diameter (mm) of the samples extracts against the bacterial strains: *Esherichia coli*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa* and *Bacillus subtilis***

**INHIBITION ZONE DIAMETER, IZD (MM) AT A CONCENTRATION OF 100 MG/ML**

<b>BACTERIAL STRAIN</b>	<b>ETHANOL EXTRACT OF RIND (EER)</b>	<b>ETHANOL EXTRACT OF SEED (EES)</b>	<b>AQUEOUS EXTRACT OF RIND (AER)</b>	<b>AQUEOUS EXTRACT OF SEED (AES)</b>	<b>MEAN SUSCEPTIBILITY TO EXTRACTS</b>
<b><i>ESHERICHIA COLI</i></b>	6.0	4.0	4.0	4.0	4.50
<b><i>KLEBSIELLA PNEUMONIA</i></b>	4.0	4.0	5.0	3.0	4.00
<b><i>PSEUDOMONAS AERUGINOSA</i></b>	5.0	3.0	4.0	5.0	4.25
<b><i>BACILLUS SUBTILIS</i></b>	4.0	4.0	3.0	3.0	3.50
<b>MEAN ACTIVITY (INCLUDING VALUES FOR <i>B. PUMILUS</i> AND <i>S. AUREUS</i>)</b>	4.0	3.3	3.3	3.0	

Simple mean of duplicate measurements.

107  
108  
109  
110  
111  
112  
113  
114

115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133

**Table 2: Activity/inhibition zone diameter (mm) of the samples extracts against the fungal strains: *Candida albican*, *Aspergillus niger*, *Penicillium chrysogenum* and *Trichosporo bePELLI***

INHIBITION ZONE DIAMETER, IZD (MM) AT A CONCENTRATION OF 100 MG/ML					
<b>FUNGAL STRAIN</b>	<b>ETHANOL EXTRACT OF RIND (EER)</b>	<b>ETHANOL EXTRACT OF SEED (EES)</b>	<b>AQUEOUS EXTRACT OF RIND (AER)</b>	<b>AQUEOUS EXTRACT OF SEED (AES)</b>	<b>MEAN SUSCEPTIBILITY TO EXTRACTS</b>
<b><i>CANDIDA ALBICAN</i></b>	4.0	2.0	3.0	3.0	3.0
<b><i>ASPERGILLUS NIGER</i></b>	3.0	1.0	2.0	2.0	2.0
<b><i>PENICILLIUM CHRYSOGENUM</i></b>	2.9	2.0	3.0	3.0	2.5
<b><i>TRICHOSPORO BEPELLI</i></b>	1.0	0.0	2.0	2.0	1.25
<b>MEAN ACTIVITY</b>	2.5	1.25	2.5	2.5	

Simple mean of duplicate measurements.

134  
135  
136  
137  
138  
139  
140  
141

For the bacterial strains, the highest mean activity (4.0) and mean susceptibility (4.50), respectively were in the EER and *Esherichia coli* whereas the least mean activity (3.0) and mean susceptibility (1.50) were in AES and *Staphylococcus aureus* (Table 1, Figures 1 and 2). The highest mean activity (2.5) and mean susceptibility (3.0), respectively for the fungal strains were in the EER, AER, AES and *Candida albican* whereas the least mean activity (1.25) and mean susceptibility (1.25) were in EES and *Trichosporo bePELLI* (Table 2).





Figure 1: Activity/inhibition zone diameter (mm) of the samples extracts at 100 mg/ml against *Bacillus pumilus*

142  
 143  
 144  
 145  
 146  
 147  
 148  
 149  
 150  
 151  
 152  
 153  
 154  
 155  
 156



**Figure 2: Activity/inhibition zone diameter (mm) of the samples extracts at 100 mg/ml against *Staphylococcus aureus***

Generally, the results revealed that the extracts were effective against the tested pathogens. The ethanol extracts of the samples however showed high antibacterial activities than the aqueous extracts, in consonance with the result of [15]. In particular, the activity of the extracts against *P. aeruginosa* were higher than that reported in Obasi *et al.* [5] for extracts and fractions of *Jatropha curcas* stem bark. The activities of the extracts against *Bacillus subtilis* and *Staphylococcus aureus* were lower than what were reported by Obasi *et al.* [2] for methanol and ethyl acetate extracts of *Samanea saman* pods at 100 mg/ml and even 50 mg/ml. The EER exhibited the highest activity against the bacterium, *Escherichia coli* (6.0) and the fungus, *Candida albican* (4.0), indicating the overriding efficacy of EER against the pathogens. Antibacterial and antifungal activities were attributed to the presence of antinutrients [16,17], warranting further studies on the antinutrient compositions of the watermelon rind and seed.

For the bacterial strains, the highest mean activity (4.0) and mean susceptibility (4.50), respectively were in the EER and *Escherichia coli* whereas the least mean activity (3.0) and mean susceptibility (1.50) were recorded in AES and *Staphylococcus aureus*. This suggests the higher potency of EER over others (EES, AES and AER) against ailments caused by bacteria, notably *Escherichia coli*. For the fungal strains, the highest mean activity (2.5) and mean susceptibility (3.0), respectively were in the EER, AER, AES and *Candida albican*, suggesting the potency of these extracts of watermelon against *Candida albican* and possibly diseases caused by *Candida albican*. The least mean activity against (1.25) and mean susceptibility to (1.25) the fungal strains were recorded in EES and *Trichosporo bepellii*. Furthermore, the least activity was recorded in EES against *Trichosporo bepellii* (0). These imply that the extract (EES) could neither combat *Trichosporo bepellii* nor serve as broad spectrum antibiotic. The activities were higher in the EE than in the AE of the rind and seed, suggesting the preference to the EE for higher pharmacologic activity. The extracts of the rind has higher antibacterial and antifungal activities than that of the seed, suggesting that the rind could be a better source for pharmacologic agents against the studied pathogens.

#### 4. CONCLUSION

The extracts (EER, AER, EES and AES) exhibited activities against the tested strains, especially *Escherichia coli* and *Candida albicans*, hence scientific basis for the utilization of the extracts in nutraceutical formulations and in ethnomedication. The study underscores the overriding potency of EER against the bacterium, *Escherichia coli* and the fungus, *Candida albicans*, and the non-susceptibility of the fungus, *Trichosporo bepellii* to EES. Further works, however are needed to validate reliability.

190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244

## REFERENCES

1. Anderson ER, Koplan IJ, Henney JE. Diagnosis and management of food borne illness. A primer for physicians. Center for Disease Control Morbidity and Mortality Weekly Report. 2001; 50(2):1-69.
2. Obasi NL, Egbuonu ACC, Ukoha PO, Ejikeme PM. Comparative Phytochemical and Antimicrobial screening of some solvent extracts of *Samanea saman* (Fabaceae or Mimosaceae) pods. African Journal of Pure and Applied Chemistry. 2010; 4(9):206-212.
3. Egbuonu ACC, Nwankwo NE. Phytochemical of some solvent fractions of petroleum ether extract of the African Mistletoe (*Loranthus micranthus* Linn) leaves and their antimicrobial activity. African Journal of Biotechnology. 2012; 11(62):12595-12599.
4. Ukoha PO, Egbuonu ACC, Obasi NL, Ejikeme PM. Tannins and other phytochemicals of the *Samanea saman* pods and their antimicrobial activities. African Journal of Pure and Applied Chemistry. 2011; 5(8):237-244.
5. Obasi NL, Ejikeme PM, Egbuonu ACC. Antimicrobial and phytochemical activity of methanolic extract and its fractions of *Jatropha curcas* Linn (Euphorbiaceae) stem bark. African Journal of Pure and Applied Chemistry. 2011; 5(5):92-96.
6. Ajayi IA, Ajibade O, Oderinde RA. Preliminary Phytochemical Analysis of Some Plant Species. Research Journal of Chemical Science. 2011; 3(12):56-82.
7. Edeoga HO, Okwu DE, Mbaebia BO. Phytochemical constituents of some Nigerian medicinal plants. African Journal of Biotechnology. 2005; 4:685-688.
8. Yamaguchi M. 2006. World Vegetables: Principles, Production and Nutritive Values. AVI Publishing Company, Westport, Connecticut, USA. 2006; Online Edition: doi: 10.1002/food.19840281005.
9. Ensminger AH, Ensminger MK.. Food for health: A nutrition encyclopedia. Choris California Pegus Press. 1986; 1061-1072.
10. Taiwo AA, Agbotoba MO, Oyedepo A, Shobo OA, Oluwadare I, Olawuni MO. Effects of drying methods on properties of watermelon (*Citrullus lanatus*) seed oil. AJFAND. 2008; 8(4):492-501.
11. Johnson JT, Iwang EU, Hemen JT, Odey MO, Effiong EE, Eteng OE. Evaluation of anti-nutritional content of watermelon *Citrullus lanatus*. Annals of Biological Research. 2012; 3(11):5145-5150
12. Oseni OA, Okoye VI. Studies of phyto-chemical and anti-oxidant properties of the fruit of watermelon (*Citrullus lanatus*), Journal of Pharmaceutical and Biomedical Science. 2013; 27(27):508-514.
13. Fila WA, Ifam EH, Johnson JT, Odey MO, Effiong EE, Dasofunjo K, Ambo EE. Comparative proximate compositions of watermelon *Citrullus Lanatus*, Squash cucurbita pepo'l and Rambutan, *Nephelium Lap-paceun*. International Journal of Science and Technology. 2013; 2(1):81-88.
14. Gin WA, Jimoh A, Abdulkareem AS, Giwa A. Production of Activated Carbon from Watermelon Peel. International Journal of Scientific & Engineering Research. 2014; 5(2):66-71.
15. Agatemor C. Antimicrobial activity of aqueous and ethanol extracts of nine Nigerian spices against four food borne bacteria. African J. Med. Sci. 2009; 255:221-224.
16. Brantner A, Males Z, Pepeljak S, Antolic A. Antimicrobial activity of *Paliurus spona-christi* Mill. (Christs thorn). J. Ethnopharmacol. 1996; 52:119-122.
17. Irobi ON, Daranola SO. Bactericidal properties of crude extracts of *Racarpu villosus*. J. Ethnopharmacol. 1994; 42:39-43.